

International Journal of Nutrition and Agriculture Research

Journal home page: www.ijnar.com



ASSOCIATION OF DIET AND PHYSICAL ACTIVITY WITH METABOLIC SYNDROME IN INDIAN ADULTS

Mahak Sharma^{*1} and Ranjana Mahna¹

^{1*}Department of Food and Nutrition, Lady Irwin College, University of Delhi, New Delhi, India.

ABSTRACT

Asians have an unusual high tendency to develop metabolic syndrome and coronary heart disease; important determinants of both these non communicable diseases are insulin resistance and clustering of other proatherogenic factors. These diseases are escalating due to marked shift in life style in Asian countries caused by economic growth, affluence, urbanization and dietary westernization. The present study was done to assess the association of metabolic syndrome, diet and physical activity in 1500 urban adults. Anthropometric measurements were assessed by standard techniques and blood pressure was measured. Blood lipid profile and blood glucose levels were assessed. Both quantitative as well as qualitative data related to diet and lifestyle was collected from the subjects. Dietary intake was determined by 24 hour recall method and food frequency questionnaire. Physical activity assessment was conducted by a suitable structured quantitative questionnaire. The results revealed that by NCEP (ATP III) criteria, 750 subjects (44.9% males and 55.1% females) with metabolic syndrome (MS) and 750 non-metabolic syndrome (NMS) subjects were identified. The main diagnostic components of MS were low HDL levels (85%) and elevated waist circumference (80%) of the MS subjects. Intake of energy and fats was higher among the MS subjects. Majority of MS subjects had low activity levels as per Physical Activity Levels (PAL) and Global guidelines for physical activity. The study concluded that unhealthy choices of foods and low physical activity are associated with a higher incidence of chronic diseases, characteristic of metabolic syndrome.

KEYWORDS

Metabolic Syndrome, Physical activity and Obesity.

Author for Correspondence:

Mahak Sharma,
Department of Food and Nutrition,
Lady Irwin College, University of Delhi,
Sikandra Road, New Delhi.

Email: mahak.sharma97@gmail.com

INTRODUCTION

Health is an important determinant of economic development growth of a country. Non-communicable diseases specially Metabolic syndrome implies a significant economic burden on health and nutrition systems, and inflict great costs on society¹.

The burden of non-communicable diseases attributed by mortality, morbidity and disability is currently

increasing and continuing to increase in developing countries, where those affected are, younger than in developed countries, and burden of 66% of these deaths occur. Unhealthy diets and lack of physical activity are among the leading causes of the all non-communicable diseases, including cardiovascular disease, diabetes and metabolic syndrome. The coexistence of high waist circumference, glucose intolerance, dyslipidemia and high blood pressure is termed as insulin resistance syndrome (IRS) or Metabolic syndrome. The clinical picture of IRS, however, may be dominated by one of its components².

Metabolic syndrome and its causes are initially mostly prevalent to economically successful groups in low and middle income countries. However, recent evidence shows that, over time, patterns of sedentary lifestyle and lack of nutrition knowledge and the metabolic syndrome associated with them spread among poor communities and contributes to social and economic inequalities. In the developing countries, even though communicable diseases and malnutrition dominate the present disease burden, but non - communicable diseases are also increasing and will grow in future scenario. The incidence of overweight and obesity are growing in developing countries, and even in low income groups in developed countries. An integrated approach to combat the causes of inadequate dietary habits and lack of physical activity would contribute to reducing the future burden of metabolic syndrome. Factors that increase the risks of metabolic syndrome include elevated consumption of energy dense, nutrient poor foods that are high in fat, sugar and salt; lack of physical activity which leads to sedentary life style at home, at work place and westernization leads to better transportation facilities.

There are a number of studies which state the relationship between metabolic syndrome, diet and physical activity in the Western countries but the researches are not as vast in the Indian context. The present study was, therefore, planned to determine the association between diet, physical activity and metabolic syndrome.

MATERIAL AND METHODS

The study was conducted in 7 hospitals with 1500 subjects. The subjects comprised of middle aged men and women visiting the OPDs for medical problems related to the components of metabolic syndrome. According to third report of National Cholesterol Education Program (NCEP (ATPIII) 2005)³, metabolic syndrome is defined as having three or more of the following abnormalities: Elevated waist circumference ≥ 40 inches for men, ≥ 35 inches for women; Elevated triglycerides ≥ 150 mg/dl; Reduced HDL Cholesterol < 40 mg/dl in men, < 50 mg/dl in women; Elevated fasting glucose ≥ 100 mg/dl; Elevated blood pressure $\geq 130/85$ mmHg. NCEP (ATP III) criteria⁵ was used to select freshly diagnosed cases of metabolic syndrome. Anthropometric measurements were done and the subjects were examined for blood pressure. Data regarding biochemical parameters viz blood glucose and lipid profile was compiled from the hospital records.

A sample size calculation was done for subjects with MS, aged 35 to 55 years ($n = 750$) at the 95% confidence level with a 5% margin of error. An equal number of non metabolic syndrome (NMS) subjects matched for age and gender were selected.

Both quantitative and qualitative data regarding diet and physical activity was collected from the subjects by using questionnaire. Physical activity was assessed by a suitable structured questionnaire developed by Bharathi, Sandhya and Vas (2000)⁴. The dietary assessment was done by 24 hour recall method and food frequency questionnaire. The data was statistically analyzed by SPSS 21 version. The project was approved by the ethics committee of Delhi University, India and all participants signed an informed consent form.

RESULTS AND DISCUSSION

Prevalence of MS Diagnostic Components

By NCEP (ATPIII) criteria, as a case subjects, 750 subjects (44.9% males and 55.1% females) were identified with metabolic syndrome (MS). An equal number (control subjects) of age and gender matched non metabolic syndrome subjects (NMS) were taken. Table No.1 illustrates the percentage prevalence of

components of metabolic syndrome in MS subjects. A large majority of MS females (89.3%) and males (85.7%) had low HDL levels - thus a major characteristic contributing to metabolic syndrome in both the genders. High waist circumference was found in MS females (82.5%) as compared to males (51.3%). Elevated blood glucose, hypertriglyceridemia and hypertension were prevalent in a fairly large number of MS males as compared to females.

Physical activity

The Global Recommendations on Physical Activity for Health aim to provide information on the relationship between physical activity and its benefits on health (i.e. the frequency, duration, intensity, type and total amount of physical activity needed for better health and prevention and treatment of NCDs and metabolic syndrome). In order to improve cardiovascular health and muscular fitness, bone and joints health and to decrease the risk of NCDs and depression and anxiety, the global recommendations on physical activity have been given by WHO (2010)¹.

In accordance with these guidelines, the physical activity of the subjects was calculated based on type of physical activity, duration and pace of physical activity reported by them.

The findings (Table No.2) revealed that a fairly large number of MS subjects (20.8% males and 30.1% females) were doing less than 150 minutes moderate/75 minutes vigorous intensity physical activity per week. The relative number of NMS subjects in this category was lesser. On the other hand, more of NMS subjects (both males 84.7% and females 82.1%) were doing physical activity more than 150 minutes of moderate/75 minutes vigorous intensity activity per week. The differences were statistically significant as per chi-square ($p=0.01$).

Table No.3 indicates PAL values, according to PAL values, a fairly large number of MS males were sedentary as compared to NMS males (65.57% vs 32.35%). Similarly, more of MS female subjects were sedentary as compared to NMS females (70.9% vs 45.27%). Odds ratio further showed a 3 times greater risk of MS with decrease in physical activity in terms of PAL values. A negative significant

association of PAL levels with metabolic syndrome was found.

Dietary data

Table No.4 shows that more of MS subjects reported preference for deep fried (86.7%) and shallow fried foods (89.9%) as compared to NMS subjects (75.6% and 82% respectively). The differences were statistically significant ($p<0.01$). A high frequency of consumption (5-6 times a week or more) of high sodium, high sugar and high fat foods was reported by a large number of subjects in both the groups. The consumption of vegetable oils rich in PUFA was reported by more of MS subjects (91.2%) as compared to NMS subjects (81.07%), whereas more of NMS subjects reported consumption of vegetable oil rich in MUFA as compared to MS subjects. The differences were statistically significant ($p<0.01$). Desi ghee was also reportedly consumed by a larger percentage of MS subjects (92.4%) as compared to NMS subjects (88.26%), but the difference was not statistically significant ($p=0.127$). Only 13% of MS subjects reported frequent change of oil used as compared to NMS subjects (30%) and the difference was significant ($p<0.01$) according to chi square.

Regarding type of milk, majority of both metabolic syndrome and non metabolic syndrome subjects reported consumption of full cream milk but the differences were not significant ($p=0.726$). A large majority of subjects in both the groups also reported consumption of sweetened beverages. However, more of MS subjects (91.2%) reported addition of sugar in the milk as compared to NMS subjects (76.54%) and this difference was statistically significant ($p<0.01$).

The findings thus denote that metabolic syndrome subjects were consuming more of fried foods, processed foods rich in sodium and fats and sugar based beverages which could be a contributory reason for abdominal obesity, high blood glucose levels and high blood pressure levels.

Energy and Nutrient Intake of Male Subjects

Figure No.1 gives the energy and nutrient intake of male subjects as percent of RDAs. In relation to RDAs, the energy intake was 84.2% for MS and 77% for NMS males, protein intake for MS males was 121.4% and 99.8% for NMS males. Thus, the

intake of energy and protein in terms of percent of RDA was higher for MS males as compared to NMS males and the difference was statistically significant ($p<0.01$). The amount of fat consumed was analyzed in terms of energy intake. Although the percent energy intake from fat was less than 30% in both cases, it was higher in MS males than NMS males (25.7% and 24.8% respectively). The percent intake of iron, calcium, thiamine, niacin, riboflavin and vitamin C in respect of RDAs was higher in MS males than that of NMS males. Differences were statistically significant for thiamine, niacin and calcium ($p<0.01$) but not for riboflavin ($p=0.115$), iron ($p=0.942$) and vitamin C consumption ($p=0.889$). The percent adequacy of vitamin A intake was low for both the groups and the difference was not statistically significant ($p=0.992$).

Energy and Nutrient Intake of Female Subjects

The intake of energy and protein with regard to RDAs (Figure No.2) was higher for MS females (92.8% and 107.8% respectively) than NMS females (68.59% and 89.5% respectively) and the difference was statistically significant ($p<0.01$). The percent of energy intake from fat was higher in MS females than NMS females (26.8% and 21.8% respectively).

The percentage intake of calcium, thiamine, riboflavin and niacin in relation to RDAs was higher in MS females as compared to NMS females. Differences were statistically significant for thiamine, niacin ($p<0.01$) but not for riboflavin ($p=0.886$) and calcium intake ($p=0.964$). Both MS and NMS females had a fairly high percentage adequacy of iron intake. The percent adequacy of vitamin A intake was low for both the groups and the difference was not statistically significant ($p=0.064$).

Nutritional Factors and Metabolic Syndrome – Results of Logistic Regression Analysis

The results of logistic regression analysis are shown in Table No.5. The data reveals that energy, fats, carbohydrates and thiamine showed a strong positive, significant association with metabolic syndrome. On the other hand, protein, fiber, iron, niacin and folic acid were found to be negatively and significantly associated. No significant association was found in case of calcium, riboflavin, vitamin A and vitamin C.

Table No.1: Percentage prevalence of metabolic syndrome diagnostic components in MS subjects

| S.No | Components of Metabolic Syndrome | MS Males (%) | MS Females (%) |
|------|----------------------------------|--------------|----------------|
| 1 | Elevated BG | 66.46 | 58.59 |
| 2 | Elevated TG | 53.70 | 44.55 |
| 3 | Low HDL | 85.7 | 89.3 |
| 4 | Elevated BP | 61.43 | 54.4 |
| 5 | Elevated WC | 51.3 | 82.5 |

LEGENDS: BG: Blood Glucose; TG: Triglycerides; HDL: High Density Lipoprotein; BP: Blood Pressure; WC: Waist Circumference.

Table No.2: Distribution of subjects as per Global guidelines for physical activity

| S.No | WHO recommendations (2010) | MS Subjects | | NMS Subjects | |
|------|--|---------------|----------------|---------------|-----------------|
| | | Males (n=103) | Females (n=93) | Males (n=150) | Females (n=145) |
| 1 | < 150 minutes moderate / 75 minutes vigorous PA per week | 21 (20.8) | 28 (30.1) | 23 (15.3) | 26 (17.9) |
| 2 | ≥ 150 minutes moderate / 75 minutes vigorous PA per week | 82 (79.2) | 65 (69.9) | 127 (84.7) | 119 (82.1) |

Table No.3: Distribution of subjects by PAL values

| S.No | Physical activity level (PAL) | Males | | Females | |
|------|-------------------------------|--------------------|------------------------|--------------------|------------------------|
| | | Metabolic syndrome | Non metabolic syndrome | Metabolic syndrome | Non metabolic syndrome |
| | | n = 337 | n = 337 | n = 413 | n = 413 |
| 1 | Sedentary (1.40-1.69) | 221 (65.57) | 109 (32.35) | 293 (70.94) | 187 (45.27) |
| 2 | Moderate (1.70-1.99) | 116 (34.43) | 228 (67.65) | 120 (29.06) | 226 (54.73) |
| 3 | Heavy (2.00-2.40) | - | - | - | - |

Chi-square (p< 0.01), Odds ratio – 3.34

Table No.4: Distribution of subjects for food preferences

| Foods Liked | MS SUBJECTS | NM S SUBJECTS |
|------------------|---------------|---------------|
| | n = 750 | n = 750 |
| Boiled | 17 (2.3) | 63 (8.4) |
| Roasted | 412 (54.9) | 403 (53.7) |
| Shallow Fried** | 674 (89.9) | 615 (82) |
| Deep Fried** | 650 (86.7) | 567 (75.6) |
| Processed | n= 750 | n=750 |
| | FOODS | |
| High Sodium | 415 (55.3) | 426 (56.8) |
| High Sugar | 300 (40.0) | 230 (30.6) |
| High Fat | 357 (47.6) | 338 (45.06) |

**Denotes significant difference (p<0.01)

| Type of Fats Consumed | MS Subjects | NMS Subjects |
|------------------------------|-------------|--------------|
| | n = 750 | n = 750 |
| Vanaspati | 5 (0.66) | 25 (3.33) |
| Desi Ghee | 693 (92.4) | 662 (88.26) |
| Vegetable oil rich in PUFA** | 684 (91.2) | 608 (81.06) |
| Vegetable oil rich in MUFA** | | |
| Mustard oil | 3 (0.4) | 29 (3.86) |
| Canola oil | 5 (0.6) | 22 (2.93) |
| Olive oil | 2 (0.3) | 9 (1.2) |
| Total | 10 (1.3) | 60 (7.99) |
| Change Of Oil | | |
| YES** | 100 (13.33) | 251 (33.46) |
| NO | 650 (86.67) | 499 (66.54) |
| How Frequently | (n = 100) | (n = 251) |
| Every month | 20 (20.0) | 61(24.31) |
| Every 2 months | 52 (52.0) | 116 (46.21) |
| Every 6 months | 27(27.0) | 74(29.48) |
| After one year | 1(1.0) | 0 (0.0) |

**Denotes significant difference (p<0.01)

| Type of Milk Consumed | MS Subjects | NMS Subjects |
|------------------------------------|-------------|--------------|
| | n = 750 | n = 750 |
| Full cream milk | 518 (69.06) | 501 (66.8) |
| Toned milk | 199 (26.55) | 211 (28.14) |
| Doubled Toned milk | 28 (3.73) | 30 (4.0) |
| Skimmed milk | 5(0.66) | 8 (1.06) |
| Consumption of Sweetened Beverages | | |
| Yes | 704(93.87) | 690 (92.0) |
| No | 46 (6.13) | 60 (8.0) |
| Consumption of Sugar With Milk | | |
| Added sugar** | 684 (91.2) | 574 (76.54) |
| No sugar | 29 (3.87) | 123 (16.4) |
| Artificial sweeteners | 37 (4.93) | 53 (7.06) |

**Denotes significant difference (p<0.01)

Table No.5: Nutritional factors that contribute to metabolic syndrome - results of logistic regression analysis

| S.No | Independent Variables | Metabolic Syndrome | |
|------|-----------------------|------------------------|----------|
| | | Regression coefficient | p- value |
| 1 | Energy** | 0.172 | <0.01 |
| 2 | Protein** | -0.720 | <0.01 |
| 3 | Fat* | 1.53 | <0.05 |
| 4 | CHO* | 0.63 | <0.05 |
| 5 | Fiber** | -0.124 | <0.01 |
| 6 | Calcium | 0.000 | 0.602 |
| 7 | Iron** | -0.139 | <0.01 |
| 8 | Thiamin** | 1.75 | <0.01 |
| 9 | Riboflavin | -0.205 | 0.709 |
| 10 | Niacin** | -0.193 | <0.01 |
| 11 | Vitamin A | 0.003 | 0.05 |
| 12 | Vitamin C | 0.003 | 0.054 |
| 13 | Folic acid** | -0.033 | <0.01 |

**Denotes significant difference (p<0.01) *Denotes significant difference (p<0.05)




Figure No.1: Energy and nutrient intake as percent of RDAs of male subjects




Figure No.2: Energy and nutrient intake as percent of RDAs of female subjects

CONCLUSION

The findings of the study revealed a strong association between metabolic syndrome, diet and physical activity. These findings concur with those of various epidemiological studies that have examined the relationship between diet, physical activity and the metabolic syndrome. The findings of the present study depict that according to PAL and Global Recommendations of physical activity given by WHO (2010)¹⁰, more of MS subjects, both males and females, were doing less physical activity as compared to NMS subjects. These findings are similar to that of a survey conducted by the Indian component of the World Health Survey (WHS)⁸, the only national level survey thus far which found that 29% of the adult physically inactive. A quarter of men (24%) and one third of women (34%) had lack of physical activity.

The study reveals that both MS males and females were having higher intake of energy, proteins and fats which could be a strong base and cause for

prevalence of obesity/overweight, hypertension, high triglycerides and diabetes. According to Cacciapuoti (2008)³, several experimental studies have shown that metabolic syndrome and low carbohydrate diets exert opposite effects on thrombosis, because these two conditions are at opposite ends of the same spectrum. Esposito et al (2007)⁴ documented nutritional factors that may increase or decrease the prevalence of the metabolic syndrome which state that beyond weight management and low caloric diets, the diet should not have more of saturated fats, trans fats, cholesterol, sodium, and simple sugars. Brunner et al (2001)² in a cross-sectional analysis of 4497 men and 1865 women aged 39-62 years showed large consumption of both PUFA and carbohydrates linked to increase waist-hip ratio, triglycerides and LDL-cholesterol. The present study shows that a high intake of energy, fats, and proteins combined with decreased physical activity is associated with occurrence of chronic diseases.

ACKNOWLEDGEMENT

I wish to express my sincere thanks and hearty gratitude to my esteemed Supervisor Dr. Ranjana Mahna for their guidance and valuable suggestions and encouragement while carrying out this whole research study.

CONFLICT OF INTEREST

We declare that we have no conflict of interest.

BIBLIOGRAPHY

1. Ramachandram A, Shenalatha C, Satyavani K. Metabolic syndrome in urban Asian Indian adults - a population study using modified ATP III criteria, *Diabetes Research and Clinical Practice*, 60(3), 2003, 199-204.
2. Cacciapuoti F. Opposite Effects of Metabolic Syndrome and Calorie Restriction on Thrombotic Disease: Head and Tail of Same Coin-Resveratrol's Role, *Metab Syndr Relat Disord*, 7(5), 2008, 397-400.
3. National Diabetes Information Clearinghouse (NDIC), NIH Publication No.09-4893 October 2008.
4. Global strategy on diet, Physical activity and health, *The World Health Report*, 2002.
5. Mishra A and Vikram N K. Insulin resistance syndrome and obesity in Asian Indians: evidence and implication, *Indian J Med Science*, 20(5), 2004, 482-491.
6. Grundy S M, Cleeman J I, Daniels S R *et al*. American Heart Association, National Heart, Lung, and Blood Institute. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement, *Circulation*, 112(17), 2005, 2735-2752.
7. Bharathi A V, Sandhya, Vaz M. The development and characteristics of a physical activity questionnaire for epidemiological studies in urban middle class Indians, *Indian J Med Res*, 111, 2000, 95-102.
8. Esposito K, Ceriello A, Giugliano D. Diet and the metabolic syndrome, *Metab Syndr Relat Disord*, 5(4), 2007, 291-6.
9. Brunner E J, Wunsch H, Marmot M G. What is an optimal diet? Relationship of macronutrient intake to obesity, glucose tolerance, lipoprotein cholesterol levels and the metabolic syndrome in the Whitehall II study, *International Journal of Obesity and Related Metabolic Disorders*, 25(1), 2001, 45-53.
10. Mohan V, Gokulakrishnan K, Deepa R *et al*. Association of physical inactivity with components of metabolic syndrome and coronary artery disease- the Chennai urban population study, *Diabetic Medicine*, 22(9), 2005, 1206-1211.

Please cite this article in press as: Mahak Sharma and Ranjana Mahna. Association of diet and physical activity with metabolic syndrome in Indian adults, *International Journal of Nutrition and Agriculture Research*, 1(1), 2014, 20-28.